

## Table of Contents

<b>Detection of <i>bla</i><sub>TEM</sub> &amp; <i>bla</i><sub>SHV</sub> genes among clinical isolates of <i>E. coli</i> from Tehran hospitals</b>	<b>I</b>
Shahcheraghi F*, Noveiri H, Nasiri S	
<b>The prevalence and molecular characterization of vancomycin resistant gram positive cocci isolated from patients in Tehran</b>	<b>II</b>
Eshraghi SS*, Talebi M, Pourshafie M. R, Salari MH	
<b>Cytolethal distending toxin (CDT) produced by <i>Campylobacter jejuni</i> and <i>Campylobacter coli</i> isolated from chickens by tissue culture method in Isfahan</b>	<b>III</b>
Havaei SA* , Pishva E, Tabibian A, Rabani M, Haghshenas F, Narimani T	
<b>Determine the inducible resistance phenotype in methicillin resistance <i>staphylococcus aureus</i> and coagulase negative staphylococci</b>	<b>IV</b>
Naderinasab M*, Yousefi F, Farshadzadeh Z, Sasan M	
<b>Isolation of carbapenem resistant <i>Acinetobacter baumannii</i> (CRAB) strains from patients and equipments of Intensive care units (ICUs) at Qazvin between 2005-2006</b>	<b>V</b>
Khosroshahi N*, Sharifi M	
<b>Frequency of enterohaemorrhagic <i>Escherichia coli</i> isolated from patients with acute diarrhea in Tabriz hospitals</b>	<b>VI</b>
Nahaei MR * , Akbari Dibavar M, Sadeghi J, Nikvash S	
<b>Improved Diagnosis of Brucella in human serum samples by double PCR assay</b>	<b>VII</b>
Hedayati MH*, Adli Moghaddam A, Norouzian D, Tabaraei B, Ahmadi H, Siadat SD	
<b>Seroepidemiology of HIV, syphilis, Hepatitis B and C in intravenous drug users at Loghman Hakim hospital</b>	<b>VIII</b>
Aminzadeh Z*, Aghazadeh Sarhangi K	