

## Table of Contents

<b>Using of multilocus enzyme electrophoresis to study the heterogenicity of clinical isolates of <i>Mycobacterium kansasii</i></b> Feizabadi MM*, Bahrmand AR, Heidarieh P, Shojaee H	<b>I</b>
<b>Comparison of Oxacillin Agar Screening and PCR Methods in Detection of Methicillin Resistance <i>Staphylococcus epidermidis</i> Strains Isolated from Blood Cultures of Children</b> Shoja S, Nahaei MR*, Farajnia S, Ahangarzadeh Rezaee M, Nikvash S	<b>II</b>
<b>PCR detection of PER &amp; VEB &amp; SHV and TEM <math>\beta</math>-lactamases in multidrug resistant <i>P. aeruginosa</i> isolated from wound infections in two hospitals of Tehran</b> Shahcheraghi F*, Nikbin VS, Shooraj F	<b>III</b>
<b>Rapid detection of <i>Pseudomonas aeruginosa</i> in clinical samples of burned patients by Fluorescent in situ hybridization(FISH)</b> Badie F*, Moosavian M	<b>IV</b>
<b>Prevalence of <i>Campylobacter jejuni</i> samples from patients referred to Semnan public health centers in 2007</b> Irajian GR, Jazayeri Moghadas A*, Beheshti AS, Salehian A, Monem M, Ghods F	<b>V</b>
<b>Molecular Detection of bacterial pathogens involved in urinary tract infection</b> Sedighian H, Pourmand MR*	<b>VI</b>
<b>Assessment of the S-layer nanostructure production under anaerobic and 5% CO<sub>2</sub> conditions in <i>Lactobacillus acidophilus</i> ATCC4356</b> Khaleghi M*, Kasra Kermanshahi R, Zarkesh SH	<b>VII</b>
<b>Comparison of the antibacterial activity of Handsept and Decosept</b> Rokoei F, Rezaei S, Tehrani M, Karbasian M, Rastegar Lari A*	<b>VIII</b>